

Accounting for running α_s for the non-singlet components of the structure functions F_1 and g_1 at small x .

B. I. Ermolaev

A.F.Ioffe Physico-Technical Institute, 194021 St.Petersburg, Russia

M. Greco

Dipartimento di Fisica and INFN, University of Rome III, Rome, Italy

S. I. Troyan

St.Petersburg Institute of Nuclear Physics, 188300 Gatchina, Russia

Infrared evolution equations incorporating the running QCD coupling are constructed and solved for the non-singlet structure functions f_{NS} . Accounting for dropped logs of x in DGLAP it leads to a scaling-like small x behaviour $f_{NS} \sim (\sqrt{Q^2}/x)^a$. In contrast to the leading logarithmic approximation, intercepts a are numbers and do not contain α_s . It is also shown that the leading logarithmic approximation may be unreliable for predicting Q^2 -dependence of the DIS structure functions in the HERA range.

Non-singlet structure functions, i.e. flavour-dependent contributions to the deep inelastic structure functions, have been the object of intensive theoretical investigation. First, they are interesting because they are experimentally measurable quantities; second, they are comparatively technically simple for analysis, and can be regarded as a starting ground for a theoretical description of DIS structure functions. In the present talk we discuss the explicit expressions³ for the non-singlet contribution f_{NS}^+ to the structure function F_1 and for the non-singlet contribution f_{NS}^- to the spin structure function g_1 at x . These expressions account for both leading (double-logarithmic) and sub-leading (single-logarithmic) contributions to all orders in QCD coupling and include the running α_s effects. Contrary to DGLAP¹ and to some other works on small x , we do not use a priori the standard parametrisation $\alpha_s = \alpha_s(Q^2)$ in our evolution eqs. Indeed it has been shown recently³ that such a dependence is a good approximation at large x but is not correct when x is small.

As we account for double-logarithmic (DL) and single-logarithmic (SL) contributions to all orders and regardless of the arguments, we cannot use the DGLAP eqs. Instead, we construct and solve two-dimensional infrared evolution equations (IREE) for f_{NS} appreciating evolution with respect to x and to Q^2 . In the context of this method, f_{NS}^\pm evolves with respect to the infrared cut-off μ in the transverse momentum space: $k_{i\perp} > \mu$ for all virtual particles. In doing so, we provide $f_{NS}^\pm(x, Q^2)$ with μ dependence too. However,

it's unavoidable when α_s is running because the standard expression

$$\alpha_s(t) = \frac{1}{b \ln(t/\Lambda_{QCD}^2)} , \quad (1)$$

is valid only when $t \gg \Lambda_{QCD}^2$ and therefore if we introduce the infrared cut-off as

$$k_{i\perp} > \mu > m_{max} \gg \Lambda_{QCD} , \quad (2)$$

with m_{max} being the mass of the heaviest involved quark, we can neglect quark masses and still do not have infrared singularities. Besides the restrictions imposed by Eq. (2) μ is not fixed, so f_{NS} can evolve with respect to μ , eventually arriving at the following expressions for the non-singlet structure functions:

$$f_{NS}^\pm = \int_{-\infty}^{\infty} \frac{d\omega}{2\pi i} C \left(\frac{1}{x} \right)^\omega \omega \exp([(1 + \lambda\omega) F_0^\pm] y) \quad (3)$$

where C is an (non-perturbative) input and F_0^\pm are the new anomalous dimensions. They account for the total resummation of the most essential at small x NLO contributions of the type $(\alpha_s/\omega^2)^n$ and $(\alpha_s/\omega)^n$ ($n = 1, \dots$) ,

$$F_0^\pm = 2 \left[\omega - \sqrt{\omega^2 - (1 + \lambda\omega)(A(\omega) + \omega D^\pm(\omega))/2\pi^2} \right] \quad (4)$$

where

$$A(\omega) = \frac{4C_F\pi}{b} \left[\frac{\eta}{\eta^2 + \pi^2} - \int_0^\infty \frac{d\rho \exp(-\rho\omega)}{(\rho + \eta)^2 + \pi^2} \right] . \quad (5)$$

and

$$D^\pm(\omega) = \frac{2C_F}{\omega b^2 N} \int_0^\infty d\rho \exp(-\rho\omega) \ln \left(\frac{\rho + \eta}{\eta} \right) \left[\frac{\rho + \eta}{(\rho + \eta)^2 + \pi^2} \mp \frac{1}{\rho + \eta} \right] . \quad (6)$$

We have used in Eqs. (5,6) the following notations: $\eta = \ln(\mu^2/\Lambda_{QCD}^2)$, $\rho = \ln(s/\mu^2)$, $\lambda = 1/2$ and the first coefficient of the β -function $b = (11N - 2n_f)/12\pi$. A corresponds to accounting for running α_s . π^2 in denominators appears due to analytical properties of $\alpha_s(t)$: it must have a non-zero imaginary part when t is time-like. D contains the signature-dependent contributions.

Expanding the resummed anomalous dimension F_0^\pm into series in $1/\omega$ we reproduce the singular in ω terms of LO and NLO DGLAP- anomalous dimensions where $\alpha_s(Q^2)C_F/2\pi$ is replaced by A .

It is shown in Refs.³ that A can be approximated by $\alpha_s(Q^2)C_F/2\pi$ only at large x . Concerning the small- x and large Q^2 asymptotics of f_{NS}^\pm , Eq. (3) reads that

$$f_{NS}^\pm \sim x^{-\omega_0^\pm} (Q^2/\mu^2)^{\omega_0^\pm/2}, \quad (7)$$

with the intercepts ω_0^\pm being the leading, i.e. the rightmost, singularities of F_0^\pm . Eqs. (3,4) read that ω_0^\pm are the rightmost roots of

$$\omega^2 - (1 + \lambda\omega)(A(\omega) + \omega D^\pm(\omega))/2\pi^2 = 0 . \quad (8)$$

Eq. (8) contains n_f, Λ_{QCD} and μ as parameters. Choosing e.g. $n_f = 3$ and $\Lambda_{QCD} = 0.1$ GeV one can solve Eq. (8) numerically and obtain ω_0^\pm as a function of μ . The solutions are given in Fig. 1. Both ω_0^+ and ω_0^- acquire imaginary parts at $\mu < 0.4$ GeV. As besides, for applicability of Eq. (1) μ must be much greater than Λ_{QCD} , we think that the region $\mu < 0.4$ GeV is beyond control of our approach. Both ω_0^+ and ω_0^- are maximal at $\mu \approx 1$ GeV and slowly decrease with μ increasing. Therefore we can estimate values of the intercepts as

$$\Omega_0^+ = 0.37, \quad \Omega_0^- = 0.4 . \quad (9)$$

It is interesting that this result was independently confirmed[†] recently by extrapolating of fits for f_3 into small x region. Eq. (9) was obtained from Eq. (8) which contains π^2 -terms. Basically, they are beyond of control of logarithmic accuracy and might be dropped. With π^2 -terms dropped, we obtain the smooth curves for ω_0^\pm depicted in Fig. 1. These curves show that π^2 -terms can be easily neglected for the values of μ greater than $\mu_0 = 5.5$ GeV. However, $\mu_0^2 = 30$ GeV² corresponds to the HERA Q^2 range. Then Eq. (9) immediately implies that, with such a big μ_0 , the logarithmic accuracy is not enough to obtain a correct Q^2 dependence in the HERA range. On the other hand, it also explains why DLA estimates $\alpha_s = \alpha_s(Q^2)$ may be correct for predicting the x dependence: indeed, in DLA where the coupling is fixed, one should use rather $\alpha_s = \alpha_s(\mu_0^2)$ than $\alpha_s = \alpha_s(Q^2)$ as taken from DGLAP, but as it happens that $\mu_0^2 = Q^2$ in the HERA range, both estimates coincide.

Acknowledgments

This work supported in part by grants INTAS-97-30494, RFBR 00-15-96610 and the EU QCDNET contract FMRX-CT98-0194.

References

1. G. Altarelli and G. Parisi, *Nucl. Phys.* **B 126** (1977) 297; V.N. Gribov and L.N. Lipatov, *Sov. J. Nucl. Phys.* **15** (1972) 438; L.N. Lipatov, *Sov. J. Nucl. Phys.* **20** (1972) 95; Yu.L. Dokshitzer, *Sov. Phys. JETP* **46** (1977) 641.
2. B.I.Ermolaev, S.I.Manayenkov and M.G.Ryskin, *Z. Phys.* **C69** (1996) 259; J.Bartels, B.I.Ermolaev and M.G.Ryskin, *Z. Phys.* **C70** (1996) 273 and **C72** (1996) 627.




Figure 1: Dependence of the intercept ω_0 on infrared cutoff μ at $\Lambda_{QCD} = 0.1$ GeV: 1– for f_1^{NS} ; 2– for g_1^{NS} ; 3– and 4– for f_1^{NS} and g_1^{NS} respectively without account of π^2 -terms.

3. B.I. Ermolaev, M. Greco and S.I. Troyan, *Nucl.Phys.* **B 571** (2000) 137 and **594** (2001) 71; hep-ph/0104082.
4. A. Kataev, G. Parente and A.V. Sidorov. *Nucl.Phys.A* 666/667 (2000) 184; *Nucl.Phys.* **B 573** (2000) 405.